

## CCAP 1686/1 *Uronema marinum*

**Note:** The culturing conditions below are not necessarily the optimal growth conditions for each strain, as much variation is found between strains, and cultures are not always kept in optimal growth conditions at CCAP for practical reasons. There may be more info in the individual strain data on the website.

**On receipt of culture:** cultures should be subcultured into fresh sterile medium as described below, ideally within a few days of receipt. If the culture vessel is very full on receipt and subculturing cannot be done immediately, we advise transferring half of the culture to a sterile container to provide air space.

**ACDP Hazard Gp:** 1 - Non pathogenic / non hazardous. Unlikely to cause human disease.

**Culture Medium:** Uronema Medium

Media recipes can be found on our website: [www.ccap.ac.uk/index.php/media-recipes/](http://www.ccap.ac.uk/index.php/media-recipes/)

**Lighting:** no light required

**Light Cycle:** -

**Temperature:** 15 degrees C

**Sub Interval:** 4 weeks

**Culture Vessel:** glass test tubes

### **Culture Method:**

Tubes containing the media are stored at 4 degrees C. One hour prior to use the required number of flasks are transferred to 15 degrees C.

To inoculate the fresh media, a dense culture is chosen from existing stocks. The state of the culture is ascertained by microscopical examination. The ciliates swim close to the meniscus.

To subculture, the inoculating tube is gently agitated to mix the cells more evenly in the medium and 1-2ml is poured aseptically into each of the new tubes.

Seal with parafilm and incubate.

**Use strict aseptic techniques throughout and if possible carry out all subculturing within a laminar flow cabinet.**